

# ALLE UNIVERD STRATES OF AMERICA

TO ALL TO WHOM THESE PRESENTS SHAM COME: Aex Gen Seed Research, III

MUCCUS, THERE HAS BEEN PRESENTED TO THE

### Secretary of Agriculture

AN APPLICATION REQUESTING A CERTIFICATE OF PROTECTION FOR AN ALLEGED DISTINCT VARIETY OF SEXUALLY REPRODUCED, OR TUBER PROPAGATED PLANT, THE NAME AND DESCRIPTION OF WHICH ARE CONTAINED IN THE APPLICATION AND EXHIBITS, A COPY OF WHICH IS HEREUNTO ANNEXED AND MADE A PART HEREOF, AND THE VARIOUS REQUIREMENTS OF LAW IN SUCH CASES MADE AND PROVIDED HAVE BEEN COMPLIED WITH, AND THE TITLE THERETO IS, FROM THE RECORDS OF THE PLANT VARIETY PROTECTION OFFICE, IN THE APPLICANT(S) INDICATED IN THE SAID COPY, AND WHEREAS, UPON DUE EXAMINATION MADE, THE SAID APPLICANT(S) IS (ARE) ADJUDGED TO BE ENTITLED TO A CERTIFICATE OF PLANT VARIETY PROTECTION UNDER THE LAW.

NOW, THEREFORE, THIS CERTIFICATE OF PLANT VARIETY PROTECTION IS TO GRANT UNTO THE SAID APPLICANT(S) AND THE SUCCESSORS, HEIRS OR ASSIGNS OF THE SAID APPLICANT(S) FOR THE TERM OF TWENTY YEARS FROM THE DATE OF THIS GRANT, SUBJECT TO THE PAYMENT OF THE REQUIRED FEES AND PERIODIC REPLENISHMENT OF VIABLE BASIC SEED OF THE VARIETY IN A PUBLIC REPOSITORY AS PROVIDED BY LAW, THE CHT TO EXCLUDE OTHERS FROM SELLING THE VARIETY, OR OFFERING IT FOR SALE, OR REPRODUCING IT, OR TING IT, OR EXPORTING IT, OR CONDITIONING IT FOR PROPAGATION, OR STOCKING IT FOR ANY OF THE URPOSES, OR USING IT IN PRODUCING A HYBRID OR DIFFERENT VARIETY THEREFROM, TO THE EXTENT BY THE PLANT VARIETY PROTECTION ACT. (84 STAT. 1542, AS AMENDED, 7 U.S.C. 2321 ET SEQ.)

FESCUE, TALL

'Regiment II'

In Testimonn Macrest, I have hereunto set my hand and caused the seal of the Plant Mariety Protection Office to be affixed at the City of Washington, D.C. this seventh day of September, in the year two thousand and seven.

Allest:

Ol-zun

Commissioner Plant Variety Protection Office Agricultural Marketing Service

REPRODUCE LOCALLY. Include form number and of	iate on all reprod	uctions			F	orm Approved - OMB No. 0581-0055				
AGRICULTURAL	NT OF AGRICUL MARKETING SEF	RVICE	The following statements are made in the Paperwork Reduction Act (PRA) of		e with the .	Privacy Act of 1974 (5 U.S.C. 552a) and				
SCIENCE AND TECHNOLOGY - F  APPLICATION FOR PLANT VA  (Instructions and information oc	RIETY PROTECT	TON CERTIFICATE	Application is required in order to dete (7 U.S.C. 2421). Information is held or	mine if a p onfidential	ant variet until certifi	y protection certificate is to be issued cate is issued (7 U.S.C. 2426).				
1. NAME OF OWNER			2. TEMPORARY DESIGNATION OR	3. VAI	RIETY NA	ME.				
NexGen Seed Research, I	LC.		ATF805	R	egiment	: II				
4. ADDRESS (Street and No., or R.F.D. No., City,	State, and ZIP Co	ode, and Country)	5. TELEPHONE (include area code)		F	OR OFFICIAL USE ONLY				
33725 Columbus St. SE			541-967-8923	PVPO NUMBER						
Albany, OR			6. FAX (include area code)	<b>│ #</b>	20	0400092				
97322			541-967-8223							
				FILING	DATE					
<ol> <li>IF THE OWNER NAMED IS NOT A "PERSON", ORGANIZATION (corporation, partnership, asso</li> </ol>		8. IF INCORPORATED, GIVE STATE OF INCORPORATION	9. DATE OF INCORPORATION		2/	2/2004				
Incorporated		Oregon	July 31, 2006		99	×4×00 T				
10. NAME AND ADDRESS OF OWNER REPRES	ENTATIVE(S) TO	 SERVE IN THIS APPLICATION, (First	person listed will receive all papers)	F	FILING	AND EXAMINATION FEES:				
Kenneth Hignight C/O				E	s 3,	652,00				
33725 Columbus St SE				R		2/2/2004				
Albany, OR 97322				E C E						
USA				1 1	1 76	· · · · · · · · · · · · · · · · · · ·				
				E	DATE	7/25/2007				
11. TELEPHONE (include area code)	12. FAX (Inclus	de area code)	13. E-MAIL		<u> </u>					
(541) 967-8923	(541) 967-8	2223								
14. CROP KIND (Common Name)	16. FAMILY N	AME (Botanical)	18. DOES THE VARIETY CONTA	AIN ANY T	RANSGE	NES? (OPTIONAL)				
Tall Fescue	Poaceae		YES NO							
15. GENUS AND SPECIES NAME OF CROP		RIETY A FIRST GENERATION HYBRI	IF SO, PLEASE GIVE THE A APPROVED PETITION TO	ISSIGNED DEREGUL	USDA-AP ATE THE	PHIS REFERENCE NUMBER FOR THE GENETICALLY MODIFIED PLANT FOR				
Festuca arundinacea	YES	I NO	COMMERICALIZATION.							
<ol> <li>CHECK APPROPRIATE BOX FOR EACH ATT (Follow instructions on reverse)</li> </ol>	ACHMENT SUBM	ITTED				HIS VARIETY BE SOLD AS A CLASS Plant Variety Protection Act)				
a.  Exhibit A. Origin and Breeding History	of the Variety		· ·			ow) 🔽 NO (If "no", go to item 23)				
b. Exhibit B. Statement of Distinctness			21. DOES THE OWNER SPECIF NUMBER OF CLASSES?	Y THAT S	EED OF T	HIS VARIETY BE LIMITED AS TO				
c. Exhibit C. Objective Description of Va	iety		☐ YES ☐ NO							
d. Z Exhibit D. Additional Description of the	Variety (Optional	)				☐ REGISTERED ☐ CERTIFIED				
e. Z Exhibit E. Statement of the Besis of the	e Owner's Owners	ship	22. DOES THE OWNER SPECIF NUMBER OF GENERATION	Y THAT S S?	EED OF T	HIS VARIETY BE LIMITED AS TO				
Voucher Sample (2,500 viable untreate verification that tissue culture will be d repository)			YES NO			540/10/400				
g.  Filing and Examination Fee (\$3,652), n	nade payable to "T	reasurer of the United	IF YES, SPECIFY THE NUMB							
States" (Mail to the Plant Variety Protect  23. HAS THE VARIETY (INCLUDING ANY HARVE)	ction Office)		FOUNDATION RE  (If additional explanation is ne	cessary, p	lease use	CERTIFIED the space indicated on the reverse.)				
FROM THIS VARIETY BEEN SOLD, DISPOSE OTHER COUNTRIES?	D OF, TRANSFER	RRED, OR USED IN THE U.S. OR				EEDER'S RIGHT OR PATENT)?				
YES NO			☐ YES ☑ NO							
IF YES, YOU MUST PROVIDE THE DATE OF FOR EACH COUNTRY AND THE CIRCUMSTA	FIRST SALE, DIS NCES. <i>(Please u</i>	POSITION, TRANSFER, OR USE is space indicated on reverse.)	IF YES, PLEASE GIVE COUN REFERENCE NUMBER. (Plea			NG OR ISSUANCE AND ASSIGNED ted on reverse.)				
25. The owners declare that a viable sample of bas a tuber propagated variety a tissue culture will t				ocordance	with such	regulations as may be applicable, or for				
The undersigned owner(s) is(are) the owner of		•		stinct, unifo	orm, and st	able as required in Section 42, and is				
entitled to protection under the provisions of Se		•								
Owner(s) is (are) informed that false represents	tion herein can jed									
SIGNATURE OF OWNER Humin			SIGNATURE OF OWNER							
NAME (Please print or type)		1	NAME (Please print or type)							
Kenneth Hignight										
CAPACITY OR TITLE  Director of Research	7	-14-0-	CAPACITY OR TITLE  Director of Research	DATE						
	l			!						

(See reverse for instructions and information collection burden statement)

#200400092

GENERAL: To be effectively filed with the Plant Variety Protection Office (PVPO), ALL of the following items must be received in the PVPO: (1) Completed application form signed by the owner; (2) completed exhibits A, B, C, E; (3) for a seed reproduced variety at least 2,500 viable untreated seeds, for a hybrid variety at least 2,500 untreated seeds of each line necessary to reproduce the variety, or for tuber reproduced varieties verification that a viable (in the sense that it will reproduce an entire plant) tissue culture will be deposited and maintained in an approved public repository; (4) check drawn on a U.S. bank for \$3,652 (\$432 filing fee and \$3,220 examination fee), payable to "Treasurer of the United States" (See Section 97.6 of the Regulations and Rules of Practice.) Partial applications will be held in the PVPO for not more than 90 days, then returned to the applicant as unfiled. Mail application and other requirements to Plant Variety Protection Office, AMS, USDA, Room 401, NAL Building, 10301 Baltimore Avenue, Beltsville, MD 20705-2351. Retain one copy for your files. All items on the face of the application are self explanatory unless noted below. Corrections on the application form and exhibits must be initiated and dated. DO NOT use masking materials to make corrections. If a certificate is allowed, you will be requested to send a check payable to "Treasurer of the United States" in the amount of \$432 for issuance of the certificate. Certificates will be issued to owner, not licensee or agent.

Plant Variety Protection Office Telephone: (301) 504-5518 FAX: (301) 504-5291

Homepage: http://www.ams.usda.gov/science/pvpo/pvpindex.htm

To avoid conflict with other variety names in use, the applicant must check the appropriate recognized authority and provide evidence that name has been cleared by the appropriate recognized authority before the Certificate of Protection is issued. For example, for agricultural and vegetable crops, contact: Seed Branch, AMS, USDA, 10301 Baltimore Avenue, Suite 401 NAL Building, Beltsville, MD 20705. Telephone: (301) 504-5682 http://www.ams.usda.gov/lsg/seed.htm.

#### ITEM

- 19a. Give:
- (1) the genealogy, including public and commercial varieties, lines, or clones used, and the breeding method;
- (2) the details of subsequent stages of selection and multiplication;
- (3) evidence of uniformity and stability; and
- (4) the type and frequency of variants during reproduction and multiplication and state how these variants may be identified
- 19b. Give a summary of the variety's distinctness. Clearly state how this application variety may be distinguished from all other varieties in the same crop. If the new variety is most similar to one variety or a group of related varieties:
  - (1) identify these varieties and state all differences objectively;
  - (2) attach statistical data for characters expressed numerically and demonstrate that these are clear differences; and
  - (3) submit, if helpful, seed and plant specimens or photographs (prints) of seed and plant comparisons which clearly indicate distinctness.
- 19c. Exhibit C forms are available from the PVPO Office for most crops; specify crop kind. Fill in Exhibit C (Objective Description of Variety) form as completely as possible to describe your variety.
- 19d. Optional additional characteristics and/or photographs. Describe any additional characteristics that cannot be accurately conveyed in Exhibit C. Use comparative varieties as is necessary to reveal more accurately the characteristics that are difficult to describe, such as plant habit, plant color, disease resistance, etc.
- 19e. Section 52(5) of the Act requires applicants to furnish a statement of the basis of the applicant's ownership. An Exhibit E form is available from the PVPO.
- 20. If "Yes" is specified (seed of this variety be sold by variety name only, as a class of certified seed), the applicant MAY NOT reverse this affirmative decision after the variety has been sold and so labeled, the decision published, or the certificate issued. However, if "No" has been specified, the applicant may change the choice. (See Regulations and Rules of Practice, Section 97.103).
- 23. See Sections 41, 42, and 43 of the Act and Section 97.5 of the regulations for eligibility requirements.
- 24. See Section 55 of the Act for instructions on claiming the benefit of an earlier filing date.
- 22. CONTINUED FROM FRONT (Please provide a statement as to the limitation and sequence of generations that may be certified.)
- 23. CONTINUED FROM FRONT (Please provide the date of first sale, disposition, transfer, or use for each country and the circumstances, if the variety (including any harvested material) or a hybrid produced from this variety has been sold, disposed of, transferred, or used in the U.S. or other countries.)
- 24. CONTINUED FROM FRONT (Please give the country, date of filing or issuance, and assigned reference number, if the variety or any component of the variety is protected by intellectual property right (Plant Breeder's Right or Patent).)

NOTES: It is the responsibility of the applicant/owner to keep the PVPO informed of any changes of address or change of ownership or assignment or owner's representative during the life of the application/certificate. The fees for filing a change of address; owner's representative; ownership or assignment; or any modification of owner's name is specified in Section 97.175 of the regulations. (See Section 101 of the Act, and Sections 97.130, 97.131, 97.175(h) of the Regulations and Rules of Practice.)

According to the Paperwork Reduction Act of 1995, an agency may not conduct or sponsor, and a person is not required to respond to a collection of information unless it displays a valid OMB control number. The valid OMB control number for this information collection is 0581-0055. The time required to complete this information collection is estimated to average 1.4 hours per response, including the time for reviewing instructions, searching existing data sources, gathering and maintaining the data needed, and completing and reviewing the collection of information.

The U.S. Department of Agriculture (USDA) prohibits discrimination in all its programs and activities on the basis of race, color, national origin, gender, religion, age, disability, sexual orientation, marital or family status, political beliefs, parental status, or protected genetic information. (Not all prohibited bases apply to all programs.) Persons with disabilities who require alternative means for communication of program information (Braille, large print, audiotape, etc.) should contact USDA's TARGET Center at 202-720-2600 (voice and TDD).

To file a complaint of discrimination, write USDA, Director, Office of Civil Rights, Room 326-W, Whitten Building, 14th and Independence Avenue, SW, Washington, DC 20250-9410 or call 202-720-5964 (voice and TDD). USDA is an equal opportunity provider and employer.

#### Exhibit A:

# Origin and Breeding History 'Regiment II' ATF805 Tall Fescue (\$7:4/2/06)

'Regiment II

The tall fescue (Festuca arundinacea) cultivar ATF805 trace

The tall fescue (Festuca arundinacea) cultivar ATF805 traces its parentage to the released cultivar Plantation. A plant selection field containing 2,491 plants of Plantation was planted in the fall of 1996. In the spring/summer of 1997 the 2,500 plants were allowed to mature allowing stem rust (Puccinia graminis) to reach full infection levels. The single plants were then rated for degree of infection. The plant selection field was flailed in early summer and the plants were rated for recovery, genetic color and crown density. One hundred plants were then selected, moved together in isolation and designated ATF592. Following harvest in 1998 a plant selection field of ATF592 was established containing 1,207 plants. In the spring of 1999 the single plants were rated for dark genetic color, crown density, number of inflorescence, level of endophyte infection (Neotyphodium coenophialum) and degree of stem rust (Puccinia graminis). Forty-two clones were selected and moved to isolation in the spring before anthesis. The 42 clones were designated ATF708, and harvested in bulk. In the fall of 1999, a single spaced-plant nursery was established of ATF708 containing 2, 200 plants. The single plants were rated for dark genetic color, crown density, number of inflorescence, level of endophyte infection (Neotyphodium coenophialum) and degree of stem rust (Puccinia graminis). Fifty clones were selected and moved to an isolation block in the spring before anthesis. The 50 clones were harvested in bulk and designated ATF805.

In the fall of 2000 a 2,097 plant breeder seed increase block was established in isolation in Albany, Oregon. ATF805 was also placed in turf trials to evaluate turf performance. The breeder seed block was harvested in bulk in 2001 and designated ATF805 (S0). A morphological nursery was established in the fall of 2001 for Plant Variety Protection (PVP) measurements.

### 2. Breeder Seed Maintenance:

A breeder seed multiplication was planted in isolation in 2000 in Albany, Oregon. Seed was harvested in bulk in 2001 and is maintained in cold storage. Seed propagation is limited to three generations, one each of foundation, registered, and certified.

## 3. Stability and Uniformity:

'Regiment II'
(ATF805) has been a stable uniform cultivar over two generations. No off-type or variant plants have been observed during the multiplication or reproduction. During the breeder seed multiplication 0.27% of the plants were removed to improve the uniformity of the population. These types were not observed during the subsequent generations. Turf plots of ATF805 have been uniform and stable.

\*Regiment II\*
Exhibit A (addendum): Statement of Stability and Uniformity for ATF805 Tall
Fescue
(N/4/06)

ATF805 has been a stable uniform cultivar over two generations. No off-type or variant plants have been observed during the multiplication or reproduction. During the breeder seed multiplication 0.27% of the plants were removed to improve the uniformity of the population. The plants that were removed showed less vigor and had poor plant health. It is not known if the lack of vigor was due to environmental factors, genetic factors, or an environment by genetic interaction. These types were not observed during the subsequent generations. Turf plots of ATF805 have been uniform and stable.

# Exhibit B: 'Regiment II' Novelty Statement of ATF805 Tall Fescue (ST:1/4/06)

The following summary outlines the distinctive characteristics of ATF805. The novelty of ATF805 is based on the unique combination of these characteristics. ATF805 is most similar to Rebel II, but may be differentiated by using the following criteria:

- 'Regiment II'
- 1) ATF805 has a later maturity (anthesis date) compared to Rebel II (tables 1A, 1B). (BT: 11/4/06)
- 2) The genetic color of ATF805 is significantly darker compared to Rebel II (tables 1A, 1B).
- 3) ATF805 has a shorter mature plant height than Rebel II (tables 1A, 1B).
- 4) The panicle length of ATF805 is shorter compared to Rebel II (tables 1A, 1B).
- 5) The flag leaf characteristics length, width, sheath length and internode length are all shorter for ATF805 compared to Rebel II (tables 1A, 1B).
- The leaf blade characteristics length and sheath length are all reduced for ATF805 compared to Rebel II (tables 1A, 1B).
- 7) The panicle length from the type most whorl to the apex is shorter for ATF805 compared to Rebel II (tables 2A, 2B).
- 8) ATF805 produces fewer florets per spikelet than Rebel II (tables 2A, 2B).
- 9) The length of the longest branch of the lowest whorl is shorter than Rebel II (tables 2A, 2B, illus. 1).
- 10) The distance between the lower most whorls is reduced compared to Rebel II (tables 2A, 2B, illus. 1).
- 11) <ATF805 has a more semi-erect growth habit compared to Rebel II (tables 3A, 3B). (657:11/4/06)

REPRODUCE LOCALLY. Include form number and date on all reproductions.

Form Approved - OMB No. 0581-0055

Public reporting burden for this collection of information is estimated to average 30 minutes per response, including the time for reviewing instructions, searching existing data sources, gathering and maintaining the data needed, and completing and reviewing the collection of information. Sand comments regarding this burden estimate or any other espect of this collection of information, including suggestions for reducing this burden, to Department of Agriculture, Clearance Officer, CIRM, AG Box 7830, Jamie L. Whitten Building, Westlington, D.C. 20250. When replying, refer to CIMB No. 0581-0055 and form number in your letter. Under the PRA of 1985, no persons are required to respond to a collection of information unless it displays a valid CIMB control number.

The U.S. Department of Agriculture (USDA) prohibits discrimination in its programs on the basis of race, color, national origin, sex, religion, age, disability, political beliefs, and marital or familial status. (Not all prohibited bases apply to all programs). Persons with disabilities who require alternative means for communication of program information (praise, large print, auditorage, etc.) should confisc the USDA Office of Communications at (202) 720-2791. To file a complaint, write the Secretary of Agriculture, U.S. Department of Agriculture, Washington, D.C. 20250, or call (202) 720-7327 (voice) or (202) 720-1127 (TDD). USDA is an equal opportunity employer.

U.S. DEPARTMENT OF AGRICULTURE AGRICULTURAL MARKETING SERVICE SCIENCE AND TECHNOLOGY PROGRAM PLANT VARIETY PROTECTION OFFICE BELTSVILLE, MD 20705 EXHIBIT C (TALL & MEADOW FESCUES)

# OBJECTIVE DESCRIPTION OF VARIETY TALL & MEADOW FESCUES

(Festuca spp.)

NAME OF APPLICANT(S) NexGen Seed Research, LLC	TEMPORARY DESIGN ATF805		TY NAME giment II
ADDRESS (Street and No., or R.F.D. No., City, State, and ZIP Code 33725 Columbus St. S. E. Albany, Oregon 97322	)	PVPO	FFICIAL USE ONLY NUMBER  0040009
Place the appropriate number that describes the varietal characteristics 089). Characteristics described, including numerical measurements, s be for SPACED PLANTS. Royal Horticultural Society or any recognit with an asterisk * are characteristics which should be recorded.	hould represent those that ar zed color fan may be used to	below. Use leading e typical for the va determine plant co	zeroes when necessary (e.g
* 1. SPECIES: (With comparison varieties, use varieties within the sp	ecies of the application vari	ety)	
$_{6}$ $_{1} = F. arundinacea (Tall)$ Turf	<u> Cypes</u>		
1 = Kentucky 31 2 = Rebel 3 = Olympic	4 = Bonanza	5 = Arid	6 = Rebel II
7 = Shortstop 8 = Silverado 9 = Rebel Jr.	10 = Mini Mustang	11 = Crewcut	12 = Bonsai
Forag	e Types		
20 = Kentucky 31 21 = Martin	22 = Forager	23 = Mozark	
24 = Kenhy 25 = AU Trium	ph 26 = Fawn	27 = Cajun	6
2 = F. pratensis (Meadow)			
30 = Admira 31 = Beaumont 32 = C	omtessa 33 = Ensign	34 = Trader	
* 2. CYTOLOGY:			
Chromosome Number			
3. ADAPTATION: (0 = Not Tested; 1 = Not Adapted; 2 = Adapted)		<del></del>	
_2_Transition Zone2_West2_Northeast	Other (Specify):		
4. MATURITY: (Date First Headed, 10% of Panicle Emergence)			
_5 Maturity Class 1 = Very early ( ) 2 = AU Triumpl	3 = Early (Fawn)	4 = K31, Kenh	y 5 = Medium (Rebel)

				· -	400092
6 = Bor	nanza 7 = La	te (Silverado)	8 = ( )	9 = Very late	
Date Headed45.75 days after A	pril 1, Locatio	onAlbany, Oreg	on, USA		
Days earlier than  Maturity same as  Days later than		son Variety			
* 5. MATURE PLANT HEIGHT from crown to top of panicle, if p			LENGTH CM: nternode subtendi	ng the flag leaf)	
_110.10_ cm Height		_2	2 .85_ cm Interne	odeLength	
_8 .53_ cm Shorter than	_6_	_1.	.73_ cm Shorter ti	han _6_ <b>\</b>	
Height same as	Comparison	Variety	Length san	ne as	Comparison Variety
cm Taller than	Comparison	— —	cm Longer	than	,
* HEIGHT AT EAR EMERGEN	CE CM: (Flag leaf height	from crown to flag	g leaf node)		
_34.83_ cm Height					
cm Shorter than	>				
Height same as	6 Comparison	Voriety			
cm Taller than	$ \begin{array}{c} -6 \\ -6 \end{array} $ Comparison	varioty			
* 6. GROWTH HABIT: (Mature P	lants)				
_7_ 1 = Prostrate (	) 3 = Ser	miprostrate ( )	5 = Ho	orizontal ( )	
7 = Semierect (R	ebel) 9 = Ere	ct (Mini Mustang)			
* 7. RHIZOMES (Psuedo):					
mm Length	$X_1 = Absent(6)$	2 = Rare (Rebel)	3 = Co	mmon ( )	
* 8. LEAF BLADE: (Tiller leaves/	turf color)				•••
*_8_ Color: 1 = Ligh	nt green ( )	3 = Medium ligh	t green (6)	5 = Green ( )	
7 = Med	lium dark green ( )	9 = Very dark gr	een ( )		
_5 Specify	rating of comparison varie	ty			
*_1_ Anthocyanin:	1 = Absent ( )	9 = Present ( )			
*_9_ Basal Hairs:	1 = Absent ( )	9 = Present ( )			
*_5_ Margins:	1 = Smooth ( )	5 = Semi-rough (	( )	9 = Rough ( )	)

8. LEAF BLADE: (continued)			<b></b> -	
*_7_Width Class: 1 =	= Very coarse ( ) $3 = C_0$	oarse ( )	5 = Medium (20)	0400092
7 =	Fine ( ) $9 = V$	ery Fine ( )		
* TILLER LEAF LENGTH CM: (First	leaf subtending the flag leaf)	* TILLE	R LEAF WIDTH MM:	
_39.85_ cm Tiller Leaf Lengt	h	_5.90_ mm	Tiller Leaf Width	
_8.75_ cm Shorter than _6	- `	mm	Narrower than	
Length same as	- Comparison Variety	Wid	th same as _6_	➤ Comparison Variety
cm Taller than	_	mm	Longer than	
FLAG LEAF LENGTH CM:		FLAG	LEAF WIDTH MM:	
_42 .53_m Flag Leaf Length		_5.15_ mm	Flag Leaf Width	
_8.37_ cm Shorter than _6_	-	_0.88_ mm	Narrower than_6_	
Length same as	Comparison Variety	Widt	h same as	Comparison Variety
cm Longer than	)	mm \	Wider than <b>J</b>	
* 9. LEAF SHEATH: (Basal Portion)				
*_1_ Anthocyanin (seedling):	1 = Absent (K31)	9 = Pre	esent ( )	
*_9_ Auricle Hairiness:	1 = Absent ( )	9 = Pre	esent (100%)	
* 10. PANICLE: (At seed maturity exce	ept where noted.)			
*_7_ Shape: 1 = Narrow-	tapering (45%) $5 = Ov$	vate ( )	7 = Oblong (55%)	9 = Other (specify)
*_7_ Type: 1 = Compac	t (45%) 5 = Int	termediate ( )	7 = Open ( 55%)	9 = Other (specify)
*_9_ Orientation: 1 =	Nodding ( )	9 = Erect ( )		
*_1_ Branch Pubescence: 1 =	Glabrous (98%)	9 = Pubescent (	)	
*_1_ Anther Color (At anthesi	s): 1 = Yellowish Green	2 = Green	3 = Bluish Green	
	4 = Purplish	5 = Reddish	6= Other (Specify)	
*_1_ Glume Color (At anthesis	s): 1 = Yellowish Green	2 = Green	3 = Bluish Green	
	4 = Purplish	5 = Reddish	6= Other (Specify)	
*_75.38_ cm Panicle Length (f	rom base to tip, if nodding, str	raighten; after anth	esis)	
_6.70_ cm Shorter than _6_				
Length same as	Comparison Variety	,		
. cm Longer than	J			

* 11. SEED: (With Lemma & Pelea)	)		
*3006 mg per 1000 se	eeds		200400092
mg Less than	<b>-</b> )		
Weight same as	— Comparison Variety		
672 mg More than	_6_ <b>)</b>		
PALEA: (Keels or Margins)	_5_ Hairs: 1 = Absent ( )	5 = Short (Missouri 96	9 = Long ( )
LEMMA:	_7_ Hairs: 1 = Absent (Kenh	5 = Several ( )	9 = Many (Missouri 96)
_6.73_ mm Lemma Length	(Mature)	_1.51_m m Lemma Width	
mm Shorter than	<b>-</b> `	mm Narrower than	<b>- \</b>
Length same as	_6_ Comparison Variety	Width same as mm Wider than	_6_ Comparison Variety
mm Longer than	_ <b>)</b>	mm Wider than	_ )
*AWNS: _9_AWN	NS: $1 = \text{Absent} ( ) 9 = \text{Pres}$	ent (Falcon) _100_% Plan	nts with awns
_0.88 mm Awn length (Of t	those present.)		
mm Shorter than _	<b>- \</b>		
Length same as	_6_ Comparison Variety		
mm Longer than _	_ }		
12. DISEASE, INSECT, AND NEM	(ATODE DEACTION: (0- Not To	ted 1- Longt Posistent 0- Man	t Degistant)
	·		
_0_ Melting-out Drechslera	a poae	_0_Blind Seed Gloeotinia temu	
_0_ Leaf Spot D. siccans		_0_ Dollar Spot <i>Lanzia, Mollero</i>	discus spp.
_0_ Net Blotch D. dictyoide	es	_0_ Stem Rust Puccinia gramin	is
_0_Brown Patch Rhizocton	nia solani	_0_ T. Blight Typhula incarnate	2
_0_ C. Leaf Spot Cercospor	ra fectucae	_0_ Pythium Blight <i>Pythium</i> spp	o.
_0_ Pink Snow Mold <i>Gerlad</i>	ichia nivalis	_0_ Powdery Mildew Erysiphe	graminis
_0_ Silver Top F. tricinctum	n, F. roseum	_0_ Crown Rust Puccinia coron	aata
_0_ Other Disease			
_0_ Other Insect			
_0_ Other Nematode			
13. ENVIRONMENTAL STRESS			
_5_ Drought Stress 1	! = Susceptible ( ) 5 = Toler	ant (6) 9 = Resistant (	( )
_5_ Shade Stress 1	1 = Susceptible ( ) 5 = Toler	ant (6) 9 = Resistant (	( )

200400092

5 Winter Stress

1 = Susceptible ( )

5 = Tolerant(6)

9 = Resistant (

14. GIVE VARIETY OR VARIETIES THAT MOST CLOSELY RESEMBLE THE APPLICATION VARIETY. For the following characteristics, indicate the degree of resemblance with the following scale:

1 = Application variety is less than comparison variety 2 = Same as 3 = More than, better, greater, darker, etc.

Character	Varieties	Rating	Character	Varieties	Rating
Leaf Width	Rebel II	1	Leaf Color	Rebel II	3
Panicle Color	Rebel II	2	Panicle Shape	Rebel II	2
Seed Size	Rebel II	3	Cold Injury	Rebel II	2
Winter Color	Rebel II	3	Heat	Rebel II	2
Disease	Rebel II	3			

<sup>\* 15.</sup> EXPERIMENTAL: Give a brief summary of the experimental design utilized to collect the data used on this form. Cultural conditions, number of plants measured and plant spacing must be specified.

A morphological nursery designated 01PVPFA was established in September 2001, in Albany, Oregon. Experimental design consisted of 20 entries; 4 replications per entry; 20 plants per replication; for a total of 80 plants per entry for tables 1A, 1B. Experimental design consisted of 20 entries; 3 replications per entry; 20 plants per replication; for a total of 60 plants per entry for tables 2 - 4. KY-31, Rebel II, Plantation and Tulsa were used as standards. Plants were established on 2.5 foot centers with a skip row between replications and between entries.

The nursery received 30 pounds of nitrogen per acre rate following establishment and 50 pounds of nitrogen per acre per year in 2002 and 2003. The fertilizer source was 15 - 15 - 15 and was applied as a split application with ½ applied in the spring and ½ in the autumn. The nursery was sprayed twice each spring, 3 weeks between applications, with Tilt (20z/acre rate), to prevent stem rust. One pound of Karmex per acre rate was applied during the late summer to prevent emergence of volunteer seedlings.

Data was analyzed using analysis of variance for a randomized complete block design. Means were calculated for each replication and then analyzed.

#### Exhibit D:

Additional Description
'RegimentII'

ATF805 Tall Fescue

ATF805 is an improved turf-type tall fescue. It has a shorter growth habit (tables 1A, 1B) than previously released tall fescue cultivars, such as KY-31, Rebel II, Tulsa and Plantation. ATF805 has a medium maturity with an anthesis date later than KY-31 and Rebel II (tables 1A, 1B). ATF805 exhibits a darker genetic color compared to KY-31, Rebel II, Tulsa, and Plantation (tables 1A, 1B). The panicle length of ATF805 is shorter compared to KY-31, Rebel II and Tulsa (tables 1A, 1B). The flag leaf characteristics; length, sheath length and internode length are all shorter for ATF805 than KY-31, Rebel II, Tulsa and Plantation (tables 1A, 1B). The flag leaf height of ATF805 is reduced compared to Plantation and Tulsa (tables 1A, 1B). The leaf blade characteristics of length and sheath length are shorter for ATF805 than KY-31, Rebel II, Plantation and Tulsa (tables 1A, 1B). The leaf blade height is reduced on ATF805 compared to KY-31 and Plantation (tables 1A, 1B). ATF805 has a shorter panicle length from the lower most whorl to the apex, compared to KY-31, Rebel II, Plantation and Tulsa (tables 2A, 2B). The whorl characteristics length of longest branch and distance between whorls is shorter for ATF805 compared to KY-31, Rebel II, Tulsa and Plantation (tables 2A, 2B, illus. 1). ATF805 has fewer plants with purple pigmentation in the panicles than Rebel II (tables 3A, 3B). ATF805 has a more erect growth habit compared to KY-31 and Tulsa, but less than Rebel II and Plantation (tables 3A, 3B). The purple pigmentation in the panicle is more frequent in ATF805 compared to KY-31, but less than Rebel II, Plantation and Tulsa (tables 3A, 3B). The presence of only one branch on the lower whorl is more frequent in ATF805 than KY-31, Rebel II and Plantation (tables 3A, 3B). ATF805 produces fewer plants with pubescence of the panicle branch than KY-31 and Plantation (tables 3A, 3B). ATF805 expresses fewer plants with dark pigmentation at the nodes compared to KY-31 (tables 4A, 4B). ATF805 has a higher seed weight per 1,000 seeds compared to KY-31, Rebel II, Plantation and Tulsa (tables 4A, 4B). ATF805 has a higher frequency of plants with semi-rough leaf blade margins compared to KY-31 and Plantation (tables 4A, 4B).

Table 1A

2002 Morphological Data

	1 4010 17	<u> </u>					<u> </u>		~							
	Cultivar	Genetic	Heading	Anthesis	Mature			Flag	Flag	Flag	Flag	Flag	Leaf	Leaf	Leaf	Leaf
		Color	Date	Date	Plant	Width	Length	Leaf	Leaf	Leaf	Leaf	Leaf	Blade	Blade	Blade	Sheath
		Scale:1-9	(days after	(days after	Height	(cm)	(cm)	Length	Width	Height	Sheath	Internode	Length	Width	Height	Length
		9999	April 1)	April 1)	(cm)			(cm)	(mm)	(cm)	Length	Length	(cm)	(mm)	(cm)	(cm)
(BT:11/4/06)	RegimentI	9-Daylost 91524) (BITGH/64 Perrequest)									(cm)	(cm)				
(81.174700)	:ATF805>	6.13	45.75	69.75	73.20	8.75	58.40	28.48	5.78	14.68	18.38	13.38	23.53	8.03	10.95	9.60
	SBL	5.71	36.25	64.75	95.18	11.03	73.18	35.28	6.70	21.93	21.83	17.25	30.05	8.85	16.48	11.73
	SBM _	5.58	39.00	65.25	88.85	11.00	67.83	32.45	6.03	21.05	20.85	17.35	35.08	8.60	15.40	11.63
	RB3	6.21	38.25	65.00	86.90	10.83	69.03	32.35	6.98	17.88	20.33	15.15	27.65	9.23	13.90	11.03
	RB2	5.90	35.50	64.50	89.40	10.40	68.85	33.45	6.78	19.45	21.00	16.68	28.25	8.93	15.03	11.25
	ATF799	5.89	43.75	66.75	83.28	10.45	62.83	29.08	6.00	20.38	19.05	17.10	26.08	8.38	14.13	10.85
	ATF800	5.61	42.50	67.25	93.05	10.78		33.65	6.00	24.75	22.25	20.48	29.40	8.30	17.05	12.10
	ATF802	5.66	40.00	65.75	91.48	11.13	72.38	34.15	5.98	18.75	22.18	16.48	27.85	8.70	13.58	10.98
	ATF704S1	5.43	38.00	65.75	97.10	11.18	70.98	35.03	6.38	25.05	22.70	20.35	29.88	8.38	17.63	12.33
	ATF803	5.59	39.50	66.75	92.48	10.60	72.73	36.48	7.45	19.63	22.10	16.65	31.48	9.35	15.10	11.83
	KY-31	3.86	35.25	65.25	122.95	11.23	83.73	48.43	7.68	37.30	32.63	26.38	44.10	11.05	31.05	18.65
	Rebel II	5.04	41.25	66.50	88.18	10.60	68.45	38.00	7.58	19.35	23.10	16.70	32.75	9.85	15.63	12.48
	Plantation	5.69	40.25	66.00	89.23	11.03	68.00	35.28	6.73	20.95	22.15	17.13	31.20	9.25	16.35	12.55
	Tulsa	5.09	40.50	66.50	97.98	11.00	72.35	37.35	6.48	24.53	23.40	19.78	32.40	8.63	18.78	13.00
	018	6.06	40.50	66.00	88.15	11.15	68.08	33.43	6.73	20.28	20.98	16.90	28.40	8.80	14.85	11.45
(80°/ <del>1</del> /11/18)	LSD(.05)	0.21	2.01	1.42	5.38	0.90	4.02	2.49	0.63	2.47	1.31	1.56	3.65	0.60	1.94	0.94
	CV _	3.31	4.32	1.82	4.89	7.02	4.83	5.97	8.27	9.33	4.96	7.18	10.05	5.81	9.83	6.54
		difference of difference of s taken in A	over two years of over one year of lbany, Oregon													

Table 1B

2003 Morphological Data

	1000						JO IVIOI	Pinolo	giçai	<u>Duta</u>						
	Cultivar		Heading	Anthesis	1 1		Panicle	Flag	Flag	Flag	Flag	Flag	Leaf	Leaf	Leaf	Leaf
		Color	Date	Date	Plant	Width	Length	Leaf	Leaf	Leaf	Leaf	Leaf	Blade	Blade	Blade	Sheath
		(Scale 1-9	(days after	(days after	Height	(cm)	(cm)	Length	Width	Height	Sheath	Internode	Length	Width	Height	Length
		9=Darkés (81:6/1/84)	April 1)	April 1)	(cm)			(cm)	(mm)	(cm)	Length	Length	(cm)	(mm)	(cm)	(cm)
'gr:।/≯/'06) `	RegimentI	per request									(cm)	(cm)				
, , , ,	ATF805>	6.34	58.50	60.83	110.10	25.75	75.38	42.53	5.15	34.83	25.80	22.85	39.85	5.90	26.30	16.43
	SBL	5.58	51.25	56.65	115.93	28.25	77.35	46.50	5.10	37.90	28.90	24.98	42.50	5.85	31.43	17.50
	SBM	5.64	57.25	59.65	111.50	27.25	71.43	43.60	5.10	38.95	26.53	24.98	40.60	6.15	31.45	16.68
	RB3	5.96	55.50	58.35	115.60	27.75	76.43	44.40	5.38	38.13	27.10	24.75	43.63	6.05	29.58	17.10
	RB2	6.06	52.00	56.48	113.75	28.00	73.23	44.68	5.00	39.88	26.63	24.70	41.95	5.38	30.45	17.50
	ATF799	5.90	60.75	61.98	107.50		67.68	38.88	4.40	38.75	23.28	25.20	38.25	5.18	28.65	16.28
	ATF800	5.76	58.50	59.83	118.05		76.53	42.60	4.85	39.60	27.40	27.35	41.48	5.40	28.30	17.78
	ATF802	_5.71	58.50	60.60	120.35		80.30	48.10	5.15	38.65	29.05	25.75	42.88	5.88	29.43	17.73
	ATF704S1	5.43	51.25	56.50	118.13		76.93	45.05	5.10	39.85	27.65	26.45	40.13	5.70	28.73	17.38
	ATF803	5.53	58.00	60.30	120.65	27.00	78.18	46.43	5.25	42.98	28.08	24.80	45.03	6.00	33.50	18.43
	KY-31	2.99	46.00	54.28	145.90	27.50	89.48	59.13	7.18	37.30	37.18	28.05	61.03	9.05	51.40	24.68
	Rebel II	5.11	57.75	59.08	118.63		82.08	50.90	6.03	36.43	28.98	24.58	48.60	6.33	28.58	18.98
	Plantation	5.71	57.75	60.28	118.68	27.75	77.60	46.80	5.68	40.40	27.85	24.75	44.43	6.78	31.53	18.33
	Tulsa	5.10	57.25	59.98	118.65	27.00	79.68	47.78	5.18	39.13	28.55	25.48	43.20	5.45	29.28	17.85
	018	6.10	59.50	60.73	108.98		70.58	44.63	5.25	38.13	26.70	23.83	42.80	6.25	29.10	16.83
BT:11/41'06)	LSD (.05)	0.25	2.47	1.15	4.51	1.77	3.91	2.13	0.78	3.57	1.44	1.16	2.42	0.92	3.55	1.06
	CV	3.90	3.74	1.65	3.25	5.47	4.33	3.90	12.72	7.51	4.34	3.86	4.67	12.97	9.61	4.99
	Cultivarium														-	

Cultivar under evaluation

<sup>■</sup> Significant difference over two years one location.
■ Significant difference over one year one location.
Measurements taken in Albany, Oregon
4 reps; 20 plants/rep = 80 data points

Table 2A

2002 Laboratory Morphological Data

_		•				~~~~	~~~		101091041	Data				
	Cultivar	Lemma	Lemma		Palea		Glume	Length of	Spikelets	Florets	Spikelet	Length of	Distance	Number of
	l	Length	Width	Awn	Length	Width	Length	Panicle from	per Panicle	per	Length	Longest	Between	Spikelets on
		(mm)	(mm)	Length	(mm)	(mm)	(mm)	Lower Most		Spikelet	(mm)	Whorl	Lower Most	the Longest
	_			(mm)				Whorl to Tip				(mm)	Whorls	Whorl
	RegimentI							(cm)					(MM) (97:43)(1871)	
(BT:11/7/76)	ATF8057	6.73	1.51	0.88	6.37	1.11	5.29	16.93	80.67	8,10	13.50	71.93	41.53	13.97
	SBL	6.85	1.46	1.06	6.45	1.15	5.36	20.53	80.00	7.10	13.00	89.57	51.30	14.83
	SBM	6.83	1.55	1.08	6.41	1.17	5.08	19.60	80.00	7.20	13.07	86.10	47.60	14.50
	RB3	6.80	1.55	0.91	6.32	1.19	5.15	18.47	76.67	7.53	13.17	76.40	46.43	13.50
	RB2	6.46	1.56	0.86	6.08	1.22	5.12	17.87	76.67	7.17	12.47	76.33	45.23	12.83
	ATF799	6.55	1.47	0.84	6.19	1.15	5.00	17.23	72.33	7.50	12.87	74.47	42.80	13.80
	ATF800	7.17	1.46	0.94	6.65	1.13	4.98	18.77	70.00	7.33	13.20	80.07	46.87	11.73
	ATF802	7.00	1.55	0.83	6.34	1.21	5.31	20.60	82.67	8.20	13.50	92.07	52.87	15.27
	ATF704S1	7.14	1.46	0.94	6.61	1.14	5.57	19.07	73.67	7.47	13.13	79.70	47.17	12.10
	ATF803	6.77	1.49	0.82	6.28	1.10	5.31	21.73	85.33	7.67	13.07	101.13	54.20	13.80
	KY-31	7.74	1.62	0.98	7.25	1.26	5.77	29.33	116.33	8.07	15.13	111.17	67.90	16.33
1	Rebel II	6.77	1.45	1.07	6.35	1.09	5.35	21.67	98.00	6.90	12.53	91.77	53.83	14.57
	Plantation	6.72	1.45	0.80	6.33	1.14	4.83	20.43	97.33	7.17	12.63	87.10	48.40	15.50
	Tulsa	6.72	1.48	0.78	6.25	1.14	4.87	21.60	93.67	7.27	12.50	89.80	50.77	13.87
	018	7.11	1.49	0.92	6.45	1.12	5.00	20.43	85.00	7.63	13.53	86.83	48.17	16.30
81:11/7/06)	LSD(.05)	0.31	0.08	0.17	0.23	0.08	0.34	2.31	11.18	0.85	0.91	13.15	5.49	2.95
	CV	3.25	3.94	12.97	2.64	5.23	4.83	8.32	9.84	8.43	5.08	11.21	8.11	15.44
	C													

Measurements taken in Albany, Oregon

Cultivar under evaluation

Significant difference over two years one location.

Significant difference over one year one location.

<sup>3</sup> reps; 20 plants/rep = 60 data points

Table 2B

2003 Laboratory Morphological Data

4.														
	Cultivar	Lemma	Lemma	Lemma		Palea	Glume	Length of	Spikelets	Florets	Spikelet	Length of	Distance	Number of
		Length	Width	Awn	Length	Width	Length	Pancile from	per Panicle	per	Length	Longest	Between	Spikelets on
		(mm)	(mm)	Length	(mm)	(mm)	(mm)	Lower Most		Spikelet	(mm)	Whorl	Lower Most	the Longest
				(mm)			Į	Whorl to Tip		,		(mm)	Whorls	Whorl
151:0/3/100	RegimentII ATF805>	<b>/</b>			1			(cm)		ļ			(mm) (m:6/i/204)	
	∕ATF805>	6.06	1.42	1.55	6.08	1.09	4.79	24.57	103.00	5.53	10.50	85.17	58.53	16.57
	SBL	7.06	1.49	1.47	6.39	1.20	5.17	26.13	92.00	5.63	11.43	101.40	67.87	16.83
	SBM	7.15	1.48	1.51	6.53	1.18	4.98	25.67	89.67	5.57	11.47	108.50	63.93	17.07
	RB3	7.16	1.45	1.48	6.50	1.22	5.07	26.23	92.67	5.27	11.27	100.10	66.67	17.93
	RB2	6.88	1.43	1.21	6.22	1.19	4.95	25.87	99.33	5.13	10.73	104.77	63.30	19.57
	ATF799	6.52	1.46	1.15	6.10	1.17	4.72	22.50	85.33	5.23	10.50	89.47	55.83	15.93
	ATF800	6.30	1.42	1.15	6.16	1.18	4.44	23.10	77.67	4.87	10.43	81.70	57.10	11.37
	ATF802	6.47	1.37	1.37	6.07	1.17	4.80	27.43	92.33	5.60	10.53	111.00	68.57	18.27
	ATF704S1	6.92	1.57	1.55	6.59	1.29	5.17	26.37	82.67	5.60	11.40	104.23	68.03	15.40
	ATF803	6.20	1.60	1.35	6.18	1.20	4.98	28.83	97.67	4.97	10.47	126.50	69.93	17.30
	KY-31	7.28	1.55	1.59	7.13	1.32	5.41	34.67	122.67	6.23	13.23	123.40	80.83	17.93
	Rebel II	6.75	1.47	1.80	6.21	1.18	5.00	29.47	108.00	4.53	10.63	117.07	71.40	18.47
	Plantation	6.41	1.45	1.31	6.14	1.19	4.40	27.30	105.33	4.80	10.13	106.87	65.43	17.37
	Tulsa	6.33	1.42	1.39	5.97	1.24	4.51	27.67	100.67	4.77	9.87	101.97	68.40	16.97
	018	6.45	1.41	1.52	6.08	1.15	4.51	25.93	92.33	5.37	10.60	96.53	63.23	16.90
BT:11/7/106)	LSD(.05)	0.70	0.10	0.27	0.31	0.10	0.37	1.81	8.48	0.47	0.79	11.74	5.46	2.80
	CV	7.74	4.86	13.52	3.64	6.31	5.60	4.93	6.51	6.53	5.28	8.27	6.05	12.18
	Cultivarium	1												

Cultivar under evaluation

3 reps; 20 plants/rep = 60 data points

Significant difference over two years one location.

Significant difference over one year one location.

Measurements taken in Albany, Oregon

## **Panicle Type Inflorescence**




Illustration 1.

Table 3A

2002 Additional Morphological Measurements of the Panicle

i		- II		Ta	A	A . 41	B 1-1-	·	A	- ·	<u> </u>	TE	- · -	- ·		<del>-</del>
	Cultivar	Growth					Panicle				Panicle	4		Panicle		5 I
	ļ	Habit at		Habit at						Orientation i				Branch	Branch	Branch
		Anthesis	Anthesis	Anthesis	Anthesis	% Purple	% Purple	% Present	% Purple	% Nodding	% Oblong	% Open	Lower	Lower	Lower	Pubescence
		% Semi-	% Horizontal	% Semi-	% Erect								Whorl	Whorl	Whorl	% Present
GT:11/7/16)	RegimentI	Prostrate		Erect									=1	=2	>3	
	ATF805>	0	20	62	18	3_	23	100	2	0	27	27	15	82	3	3
	SBL	0	20	45	35	5	20	100	2	3	60	60	30	65	5	3
	SBM	Ö	2	56	42	5	27	100	3	2	65	65	20	75	5	8
	RB3	0	28	42	30	2	18	100	0	0	72	72	30	67	3	7
	RB2	0	12	55	33	10	20	100	2	0	67	67	20	75	5	2
	ATF799	0	17	58	25	10	43	100	0	0	38	38	40	55	5	12
	ATF800	0	32	53	15	3	45	100	5	0	43	43	18	78	2	0
	ATF802	3	25	58	14	7	50	100	3	0	53	53	23	75	2	2
	ATF704S1	0	22	71	7	3	20	100	3	0	47	47	23	75	2	5
	ATF803	2	50	45	3	0	32	100	2	5	30	30	8	83	9	3
	KY-31	10	62	28	0	3	7	100	0	15	23	23	8	87	5	18
	Rebel II	3	17	48	32	3	40	100	0	0	48	48	9	80	11	3
	Plantation	0	8	64	28	5	35	100	0	0	50	50	13	85	2	12
	Tulsa	2	34	57	7	3	32	100	3	3	45	45	10	88	2	2
	018	0	5	57	38	5	25	100	0	0	45	45	40	60	0	7
	A side commercial	dat outs motio														

E Cultivar under evaluation Measurements taken in Albany, Oregon 3 reps; 20 plants/rep = 60 data points

Table 3B 2003 Additional Morphological Measurements of the Panicle

	<del></del>						9.46.11		THO THE OF						
Cultivar	Growth	Growth	Growth	Growth	Anther	Panicle	Lemma				Panicle	Panicle	Panicle	Panicle	Panicle
ţ	Habit at	Habit at			,	1	1		Orientation			Branch	Branch	Branch	Branch
	Anthesis	Anthesis	Anthesis	Anthesis	% Purple	% Purple	% Present	% Purple	% Nodding	% Oblong	% Open	Lower	Lower	Lower	Pubescence
	% Semi-	% Horizontal	% Semi-	% Erect								Whorl	Whorl	Whorl	% Present
Regiment	Prostrate		Erect		L							=1	=2	>3	
<b>∕ATF805</b> ≻	0	17	67	16	0	5	100	_ 0	0	55	55	31	59	10	2
SBL	0	22	45	33	2	12	100	3	0	65	65	34	59	7	4
SBM	0	4	55	41	2	7	100	2	0	62	62	24	68	9	2
RB3	0	25	47	28	2	3	100	0	0	66	66	39	49	13	0
RB2	0	11	57	32	2	3	100	0	0	72	72	54	44	2	1
ATF799	0	33	54	13	3	12	100	3	0	66	66	36	54	10	1
ATF800	0	41	52	7	18	0	100	3	0	70	70	21	68	11	5
ATF802	5	36	53	6	2	17	100	8	0	77	77	38	60	3	1
ATF704S1	0	25	70	5	2	5	100	_ 5	0	81	81	25	71	4	2
ATF803	6	49	43	2	2	12	100	5	0	70	70	24	71	6	1
KY-31	12	55	33	0	0	0	100	7	0	100	100	15	75	10	5
Rebel II	3	20	42	35	3	8	100	0	0	80 _	80	19	64	18	0
Plantation	_0	3	72	25	0	13	100	3	2	72	72	15	76	9	6
Tulsa	4	33	60	3	3	10	100	0	0	80	80	35	59	6	1
018	0	2	75	23	0	3	100	0	0	69	69	33	59	9	0
Cultivar un	der evaluatio	n													

Cultivar under evaluation
Measurements taken in Albany, Oregon
3 reps; 20 plants/rep = 60 data points

13

Table 4A

മാര്യ്യ (ഗ്വ് 'ഗ്?) 200% Additional Morphological Measurements

	= = p / identification into prioregrees into ded of into ite											
	Cultivar	Anthocyanin	Leaf Blade Margin	Leaf Blade Margin	Leaf Blade Margin	Leaf Blade	Leaf Sheath	Rhizomes	Lemma	Palea	Node	Seed
		Present in the	Roughness to the	Roughness to the	Roughness to the	Margin	Auricle	% Present	Hairs	Hairs	Color	Weight
	Į ,	Leaf Blade	Touch	Touch	Touch	Hairs	Hairs		% Present	% Present	% Distinct	(mg/1,000
	Regiment II	% Purple	% Smooth	% Semi-Rough	% Rough	% Present	% Present					seeds)
既川州'06)	ATE805>	0	10	12	78	100	95	0	100	100	12	3006
	SBL	0_	35	33	32	97	93	0	100	100	15	2578
	SBM	0	35	23	42	98	98	0	95	100	2	3194
	RB3	0	18	35	47	100	97	0	98	100	5	3977
	RB2	0	28	22	48	98	95	0	97	100	3	2103
	ATF799	0	25	22	53	100	97	0	100	100	7	2350
	ATF800	0	7	22	71	100	100	0	100	100	32	3080
	ATF802	0	33	24	43	100	98	0	98	100	30	_2638
	ATF704S1	0	30	32	38	100	97	0	100	100	17	2562
	ATF803	0	15	18	67	100	97	0	100	100	28	3195
	KY-31	0	58	22	18	100	95	0	100	100	30	2924
	Rebel II	0	12	15	73	100	98	0	100	100	10	2334
	Plantation	0	15	25	60	100	100	0	100	100	3	2458
	Tulsa	0	47	16	37	100	98	0	100	100	15	2347
	018	0	13	12	75	100	100	Ö	100	100	5	2338

Cultivar under evaluation
Measurements taken in Albany, Oregon
3 reps; 20 plants/rep = 60 data points

Table 4B

2003 Additional Morphological Measurements

	Cultivar —	Anthocyanin	Leaf Blade Margin	Leaf Blade Margin	Leaf Blade Margin	Leaf Blade	Leaf Sheath	Rhizomes	Lemma	Palea	Node	Seed
		Present in the	Roughness to the	Roughness to the	Roughness to the	Margin	Auricle	% Present	Hairs	Hairs	Color	Weight
		Leaf Blade	Touch	Touch	Touch	Hairs	Hairs	•	% Present	% Present	% Distinct	(mg/1,000
	Regiment 1	% Purple	% Smooth	% Semi-Rough	% Rough	% Present	% Present				}	seeds)
(BT: IV 4/ <b>8</b> 6)	ATF805>	0	82	15	3	96	90	0	99	100	0	3070
	SBL	0	85	15	0	99	90	Ō	100	100	10	2580
	SBM	0	78	17	_ 5	100	92	0	100	100	5	3190
	RB3	0	78	20	2	95	92	0	99	100	2	3973
	RB2	0	88	12	0	96	97	0	100	100	3	2109
	ATF799	0	72	18	10	92	91	0	100	100	5	2345
	ATF800	0	55	43	2	92	90	0	100	100	5	3114
	ATF802	0	83	17	0	99	90	0	100	100	18	2702
	ATF704S1	0	77	18	5	95	92	0	99	100	20	2564
	ATF803	0	70	23	7	99	95	0	100	100	10	3139
	KY-31	0	55	37	8	96	94	0	100	100	32	2937
	Rebel II	0	85	12	3	97	91	0	99	100	5	2310
'	Plantation	0	72	20	8	97	92	Ö	100	100	2	2463
	Tulsa	0	90	7	3	97	96	0	100	100	8	2352
	018	0	77	22	2	96	91	0	99	100	2	2345
	Charles and the	dor oveluation										

Cultivar under evaluation Measurements taken in Albany, Oregon 3 reps; 20 plants/rep = 60 data points

REPRODUCE LOCALLY. Include form number and edition date on all	reproductions.	ORM APPROVED - OMB No. 0581-0055					
U.S. DEPARTMENT OF AGRICULTURE AGRICULTURAL MARKETING SERVICE  EXHIBIT E  STATEMENT OF THE BASIS OF OWNERSHIP	Application is required in order to determine if a plant variety protection certificate is to be issued (7 U.S.C. 2421). The information is held confidential until the certificate is issued (7 U.S.C. 2426).						
STATEMENT OF THE BASIS OF OWNERSHIP  1. NAME OF APPLICANT(S)	2. TEMPORARY DESIGNATION	3. VARIETY NAME					
(110 1112 01 74 1 2.07411 (0)	OR EXPERIMENTAL NUMBER	o. VARIET I IVANIE					
NexGen Seed Research, LLC	ATF805	Regiment II					
4. ADDRESS (Street and No., or R.F.D. No., City, State, and ZIP, and Country)	5. TELEPHONE (Include area code)	6. FAX (include area code)					
33725 Columbus St. S. E.	(541) 967-8923	(541) 967-8223					
Albany, OR	7. PVPO NUMBER						
97322	#200400092						
8. Does the applicant own all rights to the variety? Mark an "X" in the 9. Is the applicant (individual or company) a U.S. national or a U.S. ba		السا السا					
10. Is the applicant the original owner?	NO if no, please answer <u>one</u>	of the following:					
a. If the original rights to variety were owned by individual(s), is (a YES  b. If the original rights to variety were owned by a company(ies),  YES  YES	NO If no, give name of count is (are) the original owner(s) a U.S. bas	sed company? y					
11. Additional explanation on ownership (Trace ownership from origin	al breeder to current owner. Use the re	verse for extra space if needed):					
PLEASE NOTE:							
Plant variety protection can only be afforded to the owners (not license	ees) who meet the following criteria:						
<ol> <li>If the rights to the variety are owned by the original breeder, that person must be a U.S. national, national of a UPOV member country, or national of a country which affords similar protection to nationals of the U.S. for the same genus and species.</li> </ol>							
. If the rights to the variety are owned by the company which employed the original breeder(s), the company must be U.S. based, owned by nationals of a UPOV member country, or owned by nationals of a country which affords similar protection to nationals of the U.S. for the same genus and species.							

3. If the applicant is an owner who is not the original owner, both the original owner and the applicant must meet one of the above criteria.

The original breeder/owner may be the individual or company who directed the final breeding. See Section 41(a)(2) of the Plant Variety Protection Act for definitions.

According to the Paperwork Reduction Act of 1995, an agency may not conduct or sponsor, and a person is not required to respond to a collection of information unless it displays a valid OMB control number. The valid OMB control number for this information collection is 0581-0055. The time required to complete this information collection is estimated to average 0.1 hour per response, including the time for reviewing the instructions, searching existing data sources, gathering and maintaining the data needed, and completing and reviewing the collection of information.

The U.S. Department of Agriculture (USDA) prohibits discrimination in all its programs and activities on the basis of race, color, national origin, gender, religion, age, disability, sexual orientation, marital or family status, political beliefs, parental status, or protected genetic information. (Not all prohibited bases apply to all programs.) Persons with disabilities who require alternative means for communication of program information (Braille, large print, audiotape, etc.) should contact USDA's TARGET Center at 202-720-2600 (voice and TDD).

To file a complaint of discrimination, write USDA, Director, Office of Civil Rights, Room 326-W, Whitten Building, 14th and Independence Avenue, SW, Washington, D.C. 20250-9410 or call (202) 720-5964 (voice and TDD). USDA is an equal opportunity provide and employer.